
C^* -algebras and Geometric Integration

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 03 JAN 2005		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE C*-algebras and Geometric Integration				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Heriot-Watt University Riccarton Edinburgh EH14 4AS UK				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001749, Lie Group Methods And Control Theory Workshop Held on 28 June 2004 - 1 July 2004.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 14	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Time Independent Quantum Mechanics

Let \mathcal{H} be a separable Hilbert space. Let H be a densely defined, unbounded, self adjoint operator on \mathcal{H} i.e. H may be the one dimensional Hamiltonian on $L^2(\mathbb{R})$ thus

$$Hf(x) = -\frac{1}{2}f''(x) + V(x)f(x).$$

Question: How do we compute $\sigma(H)$ the spectrum of H ?

Algorithm for computing the spectrum

- Define an appropriate one parameter family $H_\tau \in \mathcal{B}(\mathcal{H})$.
- Find a sequence of finite dimensional Hilbert spaces $\{\mathcal{H}_n\}$, $\mathcal{H}_n \subset \mathcal{H}$ with corresponding projections P_n such that $P_n \rightarrow I$ strongly and $\bigcup_{n \geq 1} \mathcal{H}_n$ is dense in \mathcal{H} with respect to the norm topology.
- Compute the eigenvalues of $A_n = P_n H_\tau|_{\mathcal{H}_n}$.

Divide the problem into two parts:

- Find the one-parameter family $H_\tau \in \mathcal{B}(\mathcal{H})$.
- Computations of the spectrum of elements in $\mathcal{B}(\mathcal{H})$.

Let $A \in \mathcal{B}(\mathcal{H})$ and let $A_n = P_n A|_{\mathcal{H}_n}$. Define

$$\Lambda = \{\lambda \in \mathbb{R} : \exists \lambda_n \in \sigma(A_n), \lambda_n \rightarrow \lambda\}.$$

For every set S of real numbers let $N_n(S)$ denote the number of eigenvalues (counting mult.) of A_n which belong to S .

Definition 1 (1) A point $\lambda \in \mathbb{R}$ is called *essential* if, for every open set U containing λ , we have

$$\lim_{n \rightarrow \infty} N_n(U) = \infty.$$

The set of essential points is denoted Λ_e

(2) $\lambda \in \mathbb{R}$ is called *transient* if there is an open set U containing λ such that

$$\sup_{n \geq 1} N_n(U) < \infty.$$

Theorem 1 (Arveson) *Let A_1, A_2, \dots be as discussed. Then $\sigma(A) \subset \Lambda$ and $\sigma_e(A) \subset \Lambda_e$.*

- Examples show that the inclusions can be proper, i.e. one may experience convergence to points not in the desired spectrum.
- We need to impose some restrictions.

Definition 2 (1) *A filtration of \mathcal{H} is a sequence $\mathcal{F} = \{\mathcal{H}_1, \mathcal{H}_2, \dots\}$ of finite dimensional subspaces of \mathcal{H} such that $\mathcal{H}_n \subset \mathcal{H}_{n+1}$ and*

$$\overline{\bigcup_{n \geq 1} \mathcal{H}_n} = \mathcal{H}$$

(2) *Let $\mathcal{F} = \{\mathcal{H}_n\}$ be a filtration of \mathcal{H} and let P_n be the projection onto \mathcal{H}_n . The degree of an operator $A \in \mathcal{B}(\mathcal{H})$ is defined by*

$$\deg(A) = \sup_{n \geq 1} \text{rank}(P_n A - A P_n).$$

Let $D(\mathcal{F})$ denote the set of all operators $A \in \mathcal{B}(\mathcal{H})$ such that

$$A = \sum_{k=1}^{\infty} A_k, \quad \deg(A_K) < \infty$$

and

$$s = \sum_{k=1}^{\infty} (1 + \deg(A_K)^{1/2}) \|A_k\| < \infty \quad (1)$$

Defining $|A|_{\mathcal{F}}$ to be the infimum of all such sums s which arise from representations of A as in (1). Then $(D(\mathcal{F}), |\cdot|_{\mathcal{F}})$ is a Banach *-algebra.

Theorem 2 (Arveson) Assume $A = A^* \in D(\mathcal{F})$. Then

(i) $\sigma_e(A) = \Lambda_e$.

(ii) Every point of Λ is either transient or essential.

Open problems

- Closing the gap $\sigma(A) \subset \Lambda$.
- Detect “false” eigenvalues.
- Definition of convergence.
- Rate of convergence.
- Error bounds in terms of n and possibly \mathcal{H}_n .
- Can the algorithm be improved?

The art of choosing H_τ

Consider the Hamiltonian H on $\mathcal{H} = L^2(\mathbb{R})$ defined by

$$H = \frac{1}{2}P^2 + v(Q),$$

where $P = -i\frac{d}{dx}$, $Q =$ multiplication by x and v is continuous. We want to find

$$P_\tau, Q_\tau \in \mathcal{B}(\mathcal{H}) \quad \text{and define} \quad H_\tau = \frac{1}{2}P_\tau^2 + v(Q_\tau).$$

Define

$$V_t f(x) = f(x - t) \quad \text{then} \quad P = \lim_{t \rightarrow 0} \frac{1}{it} (V_t - I)$$

so P is the infinitesimal generator of V_t i.e. $V_t = e^{itP}$ (Stone's Thrm).

The choice of P_τ

Possible choices of P_τ

$$(1) P_\tau = \frac{1}{i\tau}(V_\tau - I) \quad \text{or} \quad (2) P_\tau = \frac{1}{2i\tau}(V_\tau - V_{-\tau}).$$

(1) is not self-adjoint, but (2) is. Choosing $P_\tau = \frac{1}{2i\tau}(V_\tau - V_{-\tau})$ and since $V_t = e^{itP}$ we have

$$P_\tau = \frac{1}{\tau} \sin(\tau P).$$

Recalling the Spectral Mapping Theorem

$$\sigma(f(a)) = f(\sigma(a))$$

and since $f(x) = \frac{1}{\tau} \sin(\tau x)$ approximates x when τ is small suggest that P_τ could be a good choice.

The choice of Q_τ

Now P and Q satisfy the Weyl relation

$$V_t U_s = e^{ist} U_s V_t, \quad \text{where} \quad V_t = e^{itP}, U_t = e^{itQ} \quad (2)$$

which implies the “uncertainty principle”

$$PQ - QP = \frac{1}{i}I. \quad (3)$$

Now (2) cannot be achieved for P_τ, Q_τ by the Stone-von Neumann Theorem but (3) is true if

$$P = F^{-1}QF, \quad \text{where} \quad F \quad \text{is the Fourier transform.}$$

$$Q_\tau = FP_\tau F^{-1} = F \frac{1}{\tau} \sin(\tau P) F^{-1} = \frac{1}{\tau} \sin(\tau Q).$$

Simplification using representations

Theorem 3 (Arveson) *Let \mathcal{A} be the C^* -algebra generated by P_τ^2 and Q_τ and let K be a Hilbert space spanned by a bilateral orthonormal set $\{e_n : n \in \mathbb{Z}\}$. Then there is a faithful representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(K)$ such that $\pi(H_\tau)$ has the form*

$$\pi(H_\tau) = aT + bI$$

where $a = 1/8\tau^2, b = -1/4\tau^2$, and T is the tridiagonal operator

$$Te_n = e_{n-1} + 8\tau^2 v\left(\frac{1}{\tau} \sin(2n\tau)\right)e_n + e_{n+1},$$

$$n = 0, \pm 1, \pm 2, \dots$$

Open problems

- Definition of convergence.
- Rate of convergence.
- How the structure preservation affects the computational result.
- Error bounds in terms of τ .
- Final open Problem: Error bounds in terms of τ , n and \mathcal{H}_n .

Computational Chemistry

The key problem is to compute the ground state of a molecule given by

$$E_0 = \inf \{ \langle \psi, H\psi \rangle, \psi \in \mathcal{H}, \|\psi\| = 1 \}. \quad (4)$$

- Problems: E_0 may not be attained and \mathcal{H} may be huge.
- Solution: Born-Oppenheimer approximation, leads to the problem

$$\inf \{ \langle \psi, \hat{H}\psi \rangle, \psi \in \hat{\mathcal{H}}, \|\psi\| = 1 \}, \quad (5)$$

where \hat{H} and $\hat{\mathcal{H}}$ are different from (4), and (5) is attained in most cases.

- State of the art method: Born-Oppenheimer-Hartree-Fock approximation which leads to a large number of nonlinear partial differential equations.
- Possible solution: Born-Oppenheimer approximation together with the Arveson method.

Operator theory

We consider Barry Simon's 15 open problems in connection with Schrödinger operators. Given the operator on $l^2(\mathbb{Z})$ defined by

$$(h_{\alpha,\lambda,\theta}u)(n) = u(n+1) + u(n-1) + \lambda \cos(\pi\alpha + \theta)u(n),$$

where $\alpha, \lambda \in \mathbb{R}$ and $\theta \in [0, 2\pi]$

- (Problem 4)(Ten Martini) Prove that for all $\lambda \neq 0$ and all irrational α that $\sigma(h_{\alpha,\lambda,\theta})$ is a Cantor set.
- (Problem 5) Prove that for all irrational α and $\lambda = 2$ that $\sigma(h_{\alpha,\lambda,\theta})$ has measure zero.
- (Problem 6) Prove that for all irrational α and $\lambda < 2$ that the spectrum is purely continuous.